

I 蕨類植物 PTERIDOPHYTES

P1. 松葉蕨科 Psilotaceae

松葉蕨屬 *Psilotum* Sw.

松葉蕨 Pine-leaved Fern, Fork Fern

***Psilotum nudum* (L.) Griseb.**

Hab.: Herb.

Loc.: Victoria Peak, Black's Link, Tai Tam. On trees and rocks, in forests.

Dist.: S, E, SW China. Subtropical regions.

Uses: Medicinal.

P2. 石杉科 Huperziaceae

石杉屬 *Huperzia* Bernh.

華南石杉 South China Clubmoss

†***Huperzia austrosinica* Ching**

Hab.: Herb.

Loc.: Lantau Peak.

Dist.: S China.

蛇足石杉 (千層塔) Serrate Clubmoss

***Huperzia serrata* (Thunb.) Trev. (*Lycopodium serratum* Thunb.)**

Hab.: Herb.

Loc.: Tai Mo Shan, Sunset Peak. On rocks in forests and ravines.

Dist.: S, E, C & NE China. Asia.

Uses: Medicinal.

馬尾杉屬 *Phlegmariurus* (Heter) Holub

龍骨馬尾杉 Carinate Clubmoss, Keeled Clubmoss

†***Phlegmariurus carinatus* (Desv.) Ching**

Hab.: Herb.

Loc.: Cultivated.

Dist.: S China. SE Asia.

華南馬尾杉 Ford's Clubmoss

***Phlegmariurus fordii* (Baker) Ching**
(*Lycopodium fordii* Baker)

Hab.: Herb.

Loc.: Tai Mo Shan, Sunset Peak. On trunks or rocks in forests.

Dist.: Taiwan, Fujian, Guangdong, Guangxi, Yunnan. Japan, Vietnam.

P3. 石松科 Lycopodiaceae

藤石松屬 *Lycopodiastrum* Holub

藤石松 (石子藤) Climbed Clubmoss

***Lycopodiastrum casuarinoides* (Spring)**
Holub (*Lycopodium casuarinoides* Spring)

Hab.: Vine.

Loc.: Tai Mo Shan, Tai Po Kau, Ma On Shan, Lantau Island. In thickets.

Dist.: Taiwan, Fujian, Hubei, S & SW China. Tropical Asia.

Uses: Medicinal.

小石松屬 *Lycopodiella* Holub

廣東小石松 Slender Clubmoss, Carolina Clubmoss

***Lycopodiella carolinianum* (L.) Holub**
(*Lycopodium carolinianum* L.)

Hab.: Herb.

Loc.: Sha Tin, Wong Nai Chung, Mount Parker. In wet places under forest.

Dist.: Fujian, Guangdong.

燈籠草屬 *Palhinhaea* A. Franco & Vasc.

鋪地蜈蚣 (燈籠草) Nodding Clubmoss

***Palhinhaea cernua* (L.) A. Franco & Vasc.**
(*Lycopodium cernuum* L.)

Hab.: Herb.

Loc.: Common in Hong Kong. On grassy

slopes.

Dist.: E, S & SW China. Tropical Asia.

Uses: Medicinal.

P4. 卷柏科 Selaginellaceae

卷柏屬 Selaginella Beauv.

二形卷柏 Dimorphic Spikemoss

Selaginella biformis A. Br. ex Kuhn

Hab.: Herb.

Loc.: Victoria Peak, Tai Lam Chung, Tai Mo Shan, Lantau Island. In forests.

Dist.: Guangdong, Hainan, Fujian. India, SE Asia.

毛枝卷柏 Treelet Spikemoss

***Selaginella braunii** Baker

Hab.: Herb.

Loc.: Cultivated.

Dist.: E & SW China.

緣毛卷柏 Ciliated Spikemoss

Selaginella ciliaris (Retz.) Spring (*Selaginella proniflora* auct. non Baker: Check List HK Pl. 1993.)

Hab.: Herb.

Loc.: Tai Mo Shan, Lantau Island.

Dist.: Taiwan. S & SE Asia.

薄葉卷柏 Delicate Spikemoss

Selaginella delicatula (Desv. ex Poir.) Alston (*Selaginella canaliculata* Baker; *Selaginella flabellata* Spring)

Hab.: Herb.

Loc.: Common in Hong Kong. In forests.

Dist.: Areas S of Changjiang River. Vietnam, Myanmar.

深緣卷柏 Spikemoss

Selaginella doederleinii Hieron. (*Selaginella plumosa* Baker)

Hab.: Herb.

Loc.: Common in Hong Kong. In forests.

Dist.: Zhejiang, Fujian, Taiwan, Guangdong, Guangxi, Jiangxi, Hunan, Guizhou, Sichuan, Yunnan. Vietnam, Japan.

Uses: Medicinal.

兗州卷柏 Medicinal Spikemoss

Selaginella involvens (Sw.) Spring

Hab.: Herb.

Loc.: Common in Hong Kong. In thin forests.

Dist.: E, C, S & SW China. Vietnam, Myanmar, Nepal.

Uses: Medicinal.

地柏 Mat Spikemoss

***Selaginella kraussiana** (Kunze) A. Br.

Hab.: Herb.

Loc.: Cultivated.

Dist.: Native to Africa.

細葉卷柏 Slender Spikemoss

Selaginella labordei Hieron. (*Selaginella tenera* Spring)

Hab.: Herb.

Loc.: Hong Kong Island, Tai Po Kau, Lantau Island. In forests.

Dist.: S, C & SW China.

江南卷柏 Moellendorf's Spikemoss

†**Selaginella moellendorffii** Hieron.

Hab.: Herb.

Loc.: Aberdeen, Pok Fu Lam. In forests.

Dist.: Areas S of Changjiang River. Vietnam.

Uses: Medicinal.

疏葉卷柏

Selaginella remotifolia Spring (*Selaginella semicordata* Spring)

Hab.: Herb.

Loc.: Hong Kong Island, Tai Wai, Lantau Island. In wet places under forest.

Dist.: Fujian, Jiangxi, Sichuan. Tropical Asia.

卷柏 (還魂草)

†*Selaginella tamariscina* (Beauv.) Spring

Hab.: Herb.

Loc.: Common in Hong Kong. On dry cliffs.

Dist.: Widely distributed in China. E Asia.

粗葉卷柏 Rough-leaved Spikemoss

Selaginella trachyphylla A. Br. (*Selaginella atroviridis* auct. non Spring: Check List HK Pl. 1993, p. p.)

Hab.: Herb.

Loc.: Hong Kong Island. In forests.

Dist.: S China. Vietnam.

翠雲草 Blue Selaginella, Rainbow Moss, Hooked Selaginella

Selaginella uncinata (Desv.) Spring

Hab.: Herb.

Loc.: Victoria Peak. In forests.

Dist.: Zhejiang, Fujian, Taiwan, Guangdong, Guangxi, Hunan, SW China.

Uses: Medicinal, ornamental.

劍葉卷柏 Sword-leaved Spikemoss

†*Selaginella xipholepis* Bak.

Hab.: Herb.

Loc.: Victoria Peak. In forests.

P6. 木賊科 Equisetaceae

木賊屬 Equisetum L.

筆管草 (纖弱木賊) Frail Horsetail

Equisetum debile Roxb. ex Vauch.

Hab.: Herb.

Loc.: Victoria Peak, Tai Po, Wu Kau Teng. Along ditches.

Dist.: Areas S of Changjiang River. S & SE

Asia.

Uses: Medicinal.

節節草 (分枝木賊) Ramose Scouring

Rush, Many-branched Horsetail

Equisetum ramosissimum Desf.

Hab.: Herb.

Loc.: Tai Mo Shan, Tai Po Kau. On streams and wet places.

Dist.: Widely distributed in China. Temperate and subtropical regions of the old world.

Uses: Medicinal.

P9. 箭蕨科 Ophioglossaceae

箭蕨屬 Ophioglossum L.

卵葉箭蕨 Adder's Tongue-fern

Ophioglossum pedunculosum Desv.

Hab.: Herb.

Loc.: Tai Tam, Black's Link, Victoria Peak. In forests.

Dist.: Taiwan, S & SW China. Tropical Asia.

Uses: Medicinal.

心葉箭蕨 Reticulate Adder's Tongue

†*Ophioglossum reticulatum* L.

Hab.: Herb.

Loc.: Pak Ngau Shek. In forests.

Dist.: Taiwan, Fujian, Jiangxi, Guizhou, Yunnan, Sichuan. Korea, Japan, Vietnam, Malaysia, India, South America.

狹葉箭蕨

Ophioglossum thermale Kom.

(*Ophioglossum nudicaule* L. f.)

Hab.: Herb.

Loc.: Wan Chai (only one specimen collected in 1906). In forests.

Dist.: NE China, Henan, Jiangsu, Shaanxi, Jiangxi, Guangdong, Sichuan, Yunnan. Korea, Japan, Russia.

Uses: Medicinal.

P11. 蓮座蕨科 Angiopteridaceae

蓮座蕨屬 *Angiopteris* Hoffm.

福建蓮座蕨 Mules-foot Fern

**Angiopteris fokiensis* Hieron. (*Angiopteris evecta* auct. non Hoffm.: Check List HK Pl. 1993.)

Hab.: Herb.

Loc.: Mount Parker, Wong Nai Chung. In valleys.

Dist.: Fujian, Hubei, Guizhou, Guangdong, Guangxi.

Uses: Edible (young fronds), ornamental.

P13. 紫萁科 Osmundaceae

紫萁屬 *Osmunda* L.

狹葉紫萁 Narrow-leaved Osmunda

Osmunda angustifolia Ching

Hab.: Herb.

Loc.: Hong Kong Island, Sha Tin, Lantau Island. In valleys.

Dist.: Taiwan, Guangdong, Hainan.

粗齒紫萁 Gross-dentate Osmunda

†*Osmunda banksiifolia* (C. Presl) Kuhn

Hab.: Herb.

Loc.: Sha Tau Kau. In valleys.

Dist.: Guangdong, Fujian, Taiwan, Zhejiang, Japan, Philippines.

紫萁 Royal-fern

Osmunda japonica Thunb. (*Osmunda regalis* auct. non L.: Check List HK Pl. 1993.)

Hab.: Herb.

Loc.: Shing Mun. In forests.

Dist.: Areas S of Qinling Range. Korea, Japan, India.

粵紫萁 Guangdong Osmunda

Osmunda mildei C. Chr.

Hab.: Herb.

Loc.: Mount Parker, Lantau Island, Tai Mo Shan. In forests.

Dist.: Endemic to Hong Kong.

華南紫萁 Vachel's Osmunda, Vachel's Interrupted Fern

Osmunda vachellii Hook.

Hab.: Herb.

Loc.: Hong Kong Island, Tai Mo Shan, Tai Po, Lantau Island. On hillslopes or along streams.

Dist.: Fujian, Guangdong, Guangxi, Guizhou, Yunnan. Vietnam, Myanmar, India.

Uses: Ornamental.

P14. 瘤足蕨科 Plagiogyriaceae

瘤足蕨屬 *Plagiogyria* (Kunze) Mett.

華南瘤足蕨 Tumor-footed Fern

Plagiogyria tenuifolia Cop.

Hab.: Herb.

Loc.: Ma On Shan, Sunset Peak. In wet places under forest.

Dist.: Taiwan.

P15. 裹白科 Gleicheniaceae

芒萁屬 *Dicranopteris* Bernh.

鐵芒萁 Linear Forked Fern

Dicranopteris linearis (Burm. f.) Underw.

Hab.: Herb.

Loc.: Mount Parker.

Dist.: Guangdong, Hainan, Yunnan. S & SE Asia.

芒萁 Dichotomy Forked Fern

Dicranopteris pedata (Houtt.) Nakaike

[*Dicranopteris dictyotoma* Bernh;

Dicranopteris linearis auct. non (Burm. f.) Underw.: Check List HK Pl. 1993.]

Hab.: Herb.

Loc.: Common in Hong Kong. On hillslopes.
Dist.: Areas S of Changjiang River. Japan,
Vietnam, India.
Uses: Medicinal. Indicator of acidic soil.

大羽芒箕

†**Dicranopteris splendida** (Hand.-Mazz.)
Tagawa

Hab.: Herb.

Loc.: Victoria Peak, Tai Mo Shan, Pak Mong,
Man Cheung Po. In valleys.

Dist.: Hainan, Guangxi, Yunnan. Myanmar,
Vietnam.

裏白屬 **Diplopterygium** (Diels) Nakai

粵裏白

†**Diplopterygium cantonensis** (Ching) Nakai
(*Gleichenia cantonensis* Ching)

Hab.: Herb.

Loc.: Hong Kong Island, Tai Mo Shan, Fo Tan.
In thin forests.

Dist.: Guangdong.

中華裏白

†**Diplopterygium chinensis** (Ros.) DeVol

Hab.: Herb.

Loc.: Lantau Island, Tai Mo Shan. Along
ditches or in thin forests.

Dist.: Taiwan, Fujian, Guangdong, Guangxi,
Guizhou, Sichuan. Vietnam.

裏白

Diplopterygium glaucum (Thunb. ex Houtt.)
Nakai (*Hicriopteris glauca* Ching)

Hab.: Herb.

Loc.: Lion Rock, Mount Parker, Tai Mo Shan.
In thin forests.

Dist.: Areas S of Changjiang River. Japan,
India.

P17. 海金沙科 **Lygodiaceae**

海金沙屬 **Lygodium** Sw.

掌葉海金沙 Conform Climbing Fern

Lygodium conforme C. Chr. (*Lygodium dichotomum* auct. non Sw.: Check List HK Pl. 1993.)

Hab.: Climbing herb.

Loc.: Cape D'Aguilar, Tsing Yi, Castle Peak.
In thin forests.

Dist.: Guangdong, Guangxi, Hainan, Yunnan.
Vietnam.

曲軸海金沙 Flexuose Climbing Fern

Lygodium flexuosum (L.) Sw.

Hab.: Climbing herb.

Loc.: Common in Hong Kong.

Dist.: Guangdong, Hainan, Guangxi, Guizhou,
Yunnan. Vietnam, Thailand, India,
Malaysia, Australia.

海金沙 Climbing Fern

Lygodium japonicum (Thunb.) Sw.

Hab.: Climbing herb.

Loc.: Common in Hong Kong.

Dist.: Areas S of Changjiang River. Japan,
Philippines, Malaysia, India, Australia.

Uses: Medicinal.

小葉海金沙 Scansorial Climbing Fern

Lygodium scandens (L.) Sw.

Hab.: Climbing herb.

Loc.: Common in Hong Kong. In thickets.

Dist.: Taiwan, Fujian, Guangdong, Hainan,
Guangxi, Yunnan. India, Myanmar,
Malaysia.

P18. 膜蕨科 Hymenophyllaceae

假脈蕨屬 Crepidomanes Presl

南洋假脈蕨 Two-dotted Crepidomanes,

Two-punctated Crepidomanes

†**Crepidomanes bipunctatum** (Poir.) Cop.

Hab.: Herb.

Loc.: No precise locality. On moist rocks.

Dist.: Taiwan, Hainan, Guangdong, Yunnan.

Widely distributed in E Africa, S & SE Asia, Japan, Australia, New Zealand.

緣內假脈蕨

†**Crepidomanes intramarginale** (Hook. & Grev.) Cop. (*Trichomanes intramarginale* Hook. & Grev.)

Hab.: Herb.

Loc.: Mount Parker (only one specimen collected in 1906). On rocks in forest.

Dist.: Guangdong. Sri Lanka.

闊邊假脈蕨 Bristle Fern

Crepidomanes latemarginale (Eaton) Cop. (*Trichomanes latemarginale* Eaton)

Hab.: Herb.

Loc.: Mount Parker, Mount Nicholson, Tai Mo Shan. On rocks in valleys.

Dist.: Taiwan, Guangdong. Vietnam, Thailand, India.

長柄假脈蕨

Crepidomanes racemulosum (van den Bosch)

Ching (*Trichomanes orientale* auct. non C. Chr.: Check List HK Pl. 1993.)

Hab.: Herb.

Loc.: Tai Mo Shan (only one specimen collected in 1907). On trunks and rocks in forest.

Dist.: S & SW China. Vietnam, India.

團扇蕨屬 Gonocormus van den Bosch

團扇蕨 Minute Gonocormus

Gonocormus minutus van den Bosch

(*Trichomanes parvulum* auct. auct. non Baker: Check List HK Pl. 1993.)

Hab.: Herb.

Loc.: Mount Nicholson. On moist rocks.

Dist.: NE, E, S & SW China. Russia, Korea, Japan, Vietnam, Indonesia, Polynesia.

細口團扇蕨

Gonocormus nitidulus (van den Bosch) Prantl

Hab.: Herb.

Loc.: Tai Po Kau.

Dist.: Taiwan. Sri Lanka, Vietnam, Malaysia, Australia.

膜蕨屬 Hymenophyllum Sm.

華南膜蕨

***Hymenophyllum austrosinicum** Ching
(*Hymnophyllum oxyodon* auct. non Baker: Check List HK Pl. 1993.)

Hab.: Herb.

Loc.: Tai Mo Shan, Ma On Shan. On rocks in forest.

Dist.: Taiwan, Fujian, Guangdong.

露蕨屬 Mecodium Presl

露蕨 Chestnut Brown Mecodium

Mecodium badium (Hook. & Grev.) Cop.
(*Hymnophyllum javanicum* auct. non Baker: Check List HK Pl. 1993.)

Hab.: Herb.

Loc.: Lantau Island, Tai Mo Shan. On rocks in forest.

Dist.: Taiwan, Fujian, Zhejiang, Jiangxi, Anhui, Hunan, Guangdong. Korea, Japan, Vietnam, India.

小果露蕨 Small-sorus Mecodium	Hainan, Guangxi, Yunnan, Guizhou, Sichuan. Japan, Malaysia, Vietnam to India.
Mecodium microsorum (van den Bosch) Ching (<i>Hymnophyllum polyanthes</i> auct. non Sw.; <i>Hymnophyllum tenellum</i> auct. non Kuhn: Check List HK Pl. 1993.)	
Hab.: Herb.	
Loc.: Lantau Island, Tai Mo Shan, Ma On Shan. On rocks in forest.	
Dist.: E, S & SW China. Japan, Vietnam, Myanmar.	
長柄露蕨 Long-stipe Mecodium	
Mecodium osmundoides (van den Bosch) Ching	
Hab.: Herb.	
Loc.: Lantau Island, Tai Mo Shan, Pat Sin Leng.	
Dist.: S & SW China. Indochina to India.	
長筒蕨屬 <i>Selenodesmium</i> (Prantl) Cop.	
華南長筒蕨 Siamense <i>Selenodesmium</i>	
Selenodesmium siamense (Christ) Ching & C. H. Wang (<i>Trichomanes obscurum</i> auct. non Blume, <i>Trichomanes rigidum</i> auct. non Sw.: Check List HK Pl. 1993.)	
Hab.: Herb.	
Loc.: Hong Kong Island, Tai Mo Shan, Kowloon, Tai Po Kau, Sham Chung. On rocks in forest.	
Dist.: Guangdong, Hainan, Guangxi. Vietnam, Thailand.	
瓶蕨屬 <i>Trichomanes</i> L.	
瓶蕨 Bottle-fern	
Trichomanes auriculatum Blume	
Hab.: Herb.	
Loc.: Tai Mo Shan (only one specimen collected in 1907). On trunks along streams.	
Dist.: Zhejiang, Taiwan, Jiangxi, Guangdong,	
華東瓶蕨 Oriental Bottle-fern	
*Trichomanes orientale C. Chr.	
Hab.: Herb.	
Loc.: Hong Kong. On rocks along streams.	
Dist.: E, S & SW China. Japan, Korea.	
P19. 蚌殼蕨科 <i>Dicksoniaceae</i>	
金毛狗屬 <i>Cibotium</i> Kaulf.	
金毛狗 Lamb of Tartary	
Cibotium barometz (L.) J. Sm.	
Hab.: Tree-like fern.	
Loc.: Common in Hong Kong. In forests.	
Dist.: S, SW & E China. Japan, Vietnam, Myanmar, India, Malaysia.	
Uses: Medicinal, edible. Indicator of acid soil.	
P20. 條欓科 <i>Cyatheaceae</i>	
木桫欓屬 <i>Alsophila</i> R. Br.	
刺桫欓 Spiny Tree-fern	
Alsophila spinulosa (Wall. ex Hook.) Tryon (<i>Cyathea spinulosa</i> Wall. ex Hook.)	
Hab.: Tree-like fern.	
Loc.: Hong Kong Island, Lantau Island, Tai Mo Shan, Tai Lam. In valleys.	
Dist.: Taiwan, Fujian, Jiangxi, Guangdong, Guangxi, Guizhou, Sichuan.	
黑桫欓屬 <i>Gymnosphaera</i> Blume	
細齒黑桫欓 Toothed Black Tree-fern	
†Gymnosphaera hancockii (Cop.) Ching [<i>Cyathea hancockii</i> Cop.; <i>Gymnosphaera denticulata</i> (Baker) Cop.]	
Hab.: Tree-like fern.	
Loc.: Tai Mo Shan, Tung Chung. In forest of valleys.	

Dist.: Taiwan, Fujian, Jiangxi, Guangdong, Japan.

黑桫欓 Black Tree Fern

Gymnosphaera podophylla (Hook.) Cop. (*Cyathea podophylla* Cop.)

Hab.: Tree-like fern.

Loc.: Shek O, Clear Water Bay, Tseng Lan Shue. In valleys.

Dist.: Zhejiang, Fujian, Taiwan, Guangdong, Guangxi, Yunnan. Vietnam.

白桫欓屬 Sphaeropteris Bernh.

筆筒樹 Scaly Tree-fern

†***Sphaeropteris lepifera** (J. Sm. ex Hook.) Tryon (*Cyathea lepifera* Cop.)

Hab.: Tree-like fern.

Loc.: Victoria Peak (naturalized). In forests of valleys.

Dist.: Taiwan. Native to Philippines.

Uses: Ornamental.

P22. 碗蕨科 Dennstaedtiaceae

鱗蓋蕨屬 Microlepia Presl

華南鱗蓋蕨 Scaly-fern

Microlepia hancei Prantl

Hab.: Herb.

Loc.: Hong Kong Island, Sai Kung. In forests.

Dist.: Taiwan, Guangdong, Guangxi. Vietnam.

虎克鱗蓋蕨 Hooker's Scaly-fern

Microlepia hookeriana (Wall. ex Hook.) Presl

Hab.: Herb.

Loc.: Pok Fu Lam, Mount Violet, Tai Mo Shan, Ma On Shan. In forests.

Dist.: Taiwan, Fujian, Guangdong, Guangxi, Yunnan. Tropical Asia.

邊緣鱗蓋蕨 Marginal Scaly-fern

Microlepia marginata (Houtt.) C. Chr.

Hab.: Herb.

Loc.: Aberdeen, Tai Mo Shan. In forests.

Dist.: Areas S of Changjiang River. Japan, Vietnam.

粗毛鱗蓋蕨 Hairy Scaly-fern

Microlepia strigosa (Thunb.) Presl

Hab.: Herb.

Loc.: Wong Chuk Hang. In forests.

Dist.: SE coastal provinces and regions of China, Sichuan, Yunnan. Japan, Philippines, Malaysia.

P23. 鱗始蕨科 Lindsaeaceae

鱗始蕨屬 Lindsaea Dry.

劍葉鱗始蕨 (雙唇蕨) Sword-leaved Lindsaea

Lindsaea ensifolia Sw. (*Schizoloma ensifolium* J. Sm.)

Hab.: Herb.

Loc.: Common in Hong Kong. Along roadside on slopes.

Dist.: Taiwan, Fujian, Guangdong, Yunnan. Tropical Asia to SW Africa.

異葉鱗始蕨 (異葉雙唇蕨)

Lindsaea heterophylla Dry. (*Schizoloma heterophyllum* J. Sm.)

Hab.: Herb.

Loc.: Common in Hong Kong. On hills.

Dist.: Taiwan, Fujian, Guangdong, Guangxi, Yunnan. Philippines, Vietnam, Malaysia, Myanmar, India, Sri Lanka.

團葉鱗始蕨 Orbicular Lindsaea

Lindsaea orbiculata (Lam.) Mett. ex Kuhn

Hab.: Herb.

Loc.: Common in Hong Kong. In forests.

Dist.: Taiwan, Fujian, Guangdong, Guangxi, Guizhou, Sichuan. Asia, Tropical Australia.

烏蕨屬 *Stenoloma* Fée

闊片烏蕨 Bifloralet Wedgelet Fern
**Stenoloma biflorum* (Kaulf.) Ching
[*Sphenomeris chinensis* (L.) Maxon var.
littorale auct. non (Tagawa) Ohwi
Check List HK Pl. 1993.]

Hab.: Herb.
Loc.: Stanley, Kwo Chau Islands. On rocks along banks.
Dist.: Taiwan, Fujian, Guangdong. S Japan, Philippines.

烏蕨 (烏韭) Fairy Fern, Common Wedgelet Fern

Stenoloma chusanum (L.) Ching
[*Odontosoria chinensis* (L.) J. Sm.]

Hab.: Herb.
Loc.: Common in Hong Kong. In thickets.
Dist.: Areas S of Changjiang River. Japan, Philippines to Polynesia, Madagascar.
Uses: Medicinal.

P25. 姫蕨科 *Hypolepidaceae*

姬蕨屬 *Hypolepis* Bernh.

姬蕨 Downy Ground Fern, Punctate Flakelet Fern

Hypolepis punctata (Thunb.) Mett.

Hab.: Herb.
Loc.: Tai Mo Shan (only one specimen collected in 1907). In forests.
Dist.: Taiwan, Fujian, Jiangxi, Zhejiang, Anhui, Guangdong, Yunnan, Sichuan, Guizhou. Tropical and subtropical regions of Asia.

P26. 蕨科 *Pteridaceae*

蕨屬 *Pteridium* Scop.

蕨 Bracken Fern

**Pteridium aquilinum* var. *latiusculum* (Desv.) Underw. [*Pteridium aquilinum* auct. non (L.) Kuhn: Check List HK Pl.

1993.]

Hab.: Herb.

Loc.: Common in Hong Kong. In forest edges or on slopes.

Dist.: Widely distributed in China. Tropical & temperate regions.

Uses: Medicinal, edible.

P27. 凤尾蕨科 *Pteridaceae*

栗蕨屬 *Histiopteris* (Agardh) J. Sm.

栗蕨 Incised Histiopteris

Histiopteris incisa (Thunb.) J. Sm.

Hab.: Herb.

Loc.: Ma On Shan (near Wu Kau Sha, only one specimen collected in 1974). In thin forests.

Dist.: Taiwan, Fujian, Guangdong, Guangxi, Yunnan. Japan, Vietnam, Malaysia, Africa, Tropical America.

鳳尾蕨屬 *Pteris* L.

狹眼鳳尾蕨 *Biauriculae Brake*

Pteris biaurita L.

Hab.: Herb.

Loc.: Common in Hong Kong. In forests.

Dist.: Taiwan, Hainan, Guangdong, Guangxi, Yunnan. Tropical region.

條紋鳳尾蕨

Pteris cadieri Christ (*Pteris dimorpha* Cop.)

Hab.: Herb.

Loc.: Fanling (only one specimen collected in 1911). In forests.

Dist.: Taiwan, Fujian, Guangdong, Guangxi, Guizhou, Yunnan. Japan, Vietnam.

岩鳳尾蕨

Pteris deltodon Baker

Hab.: Herb.

Loc.: Ma On Shan (only one specimen collected in 1975). On limestone cliffs.

Dist.: Taiwan, Guangxi, Guizhou, Sichuan, Yunnan. Vietnam, Japan.

刺齒鳳尾蕨 Disparate Brake

†***Pteris dispar** Kunze [*Pteris semipinnata* var. *dispar* (Kunze) Hook. & Baker]

Hab.: Herb.

Loc.: Common in Hong Kong. In thin forests.

Dist.: Areas S of Changjiang River. Korea, Japan, Philippines, Malaysia, Vietnam.

劍葉鳳尾蕨 Sword Brake

Pteris ensiformis Burm.

Hab.: Herb.

Loc.: Common in Hong Kong. In forests.

Dist.: E, S & SW China. Japan, Vietnam, Myanmar, India, Sri Lanka, Malaysia to Polynesia.

Uses: Medicinal, ornamental.

金釵鳳尾蕨 Faurie's Brake

Pteris fauriei Hieron. (*Pteris quodriaaurita* auct. non Retz.: Check List HK Pl. 1993.)

Hab.: Herb.

Loc.: Mount Parker, Lamma Island. In forests.

Dist.: Zhejiang, Fujian, Taiwan, Jiangxi, Hunan, Guangdong, Guangxi, Yunnan. Japan, Vietnam.

疏裂鳳尾蕨

***Pteris finotii** Christ (*Pteris excelsa* auct. non Gaudich.: Check List HK Pl. 1993.)

Hab.: Herb.

Loc.: Nam Chung, Ng Tung Chai. In forests.

Dist.: Guangdong, Hainan, Yunnan. Vietnam.

林下鳳尾蕨 Greville's Brake

Pteris grevilleana Wall. ex Agardh

Hab.: Herb.

Loc.: Pak Tai To Yan, Ma On Shan. In forests.

Dist.: Taiwan, Guangdong, Hainan, Guangxi, Yunnan. Japan, Vietnam, Thailand, India, Nepal, Malaysia.

全緣鳳尾蕨 Conspicuous Brake

Pteris insignis Mett. ex Kuhn

Hab.: Herb.

Loc.: Tai Mo Shan, Ma On Shan, Sunset Peak, Kai Kung Leng. In forests of valleys.

Dist.: Areas S of Changjiang River. Vietnam, Malaysia.

線羽鳳尾蕨 Lineate Brake

†***Pteris linearis** Poir.

Hab.: Herb.

Loc.: Shek O. In forests.

Dist.: Taiwan, Guangdong, Hainan, Guangxi, Guizhou, Yunnan. Tropical Asia, Madagascar.

井欄邊草 Chinese Brake

Pteris multifida Poir.

Hab.: Herb.

Loc.: Common in Hong Kong.

Dist.: Areas S of Qinling Range. Japan, Philippines, Vietnam.

Uses: Medicinal.

斜羽鳳尾蕨 Oblique Pinna Brake

†**Pteris oshimensis** Hieron.

Hab.: Herb.

Loc.: Tai A Chau. In forests.

Dist.: Guangdong, Guangxi, Yunnan. Vietnam, Japan.

半邊旗 Semi-pinnated Brake

Pteris semipinnata L.

Hab.: Herb.

Loc.: Common in Hong Kong. In forests.

Dist.: Areas S of Changjiang River. Japan,
Vietnam, Malaysia, Sri Lanka, India.
Uses: Medicinal.

蜈蚣草 Ladder Brake

Pteris vittata L. (*Pteris longifolia* auct. non L.).
Check List HK Pl. 1993.)

Hab.: Herb.

Loc.: Common in Hong Kong. On calcareous
soil.

Dist.: Areas S of Qinling Range. Tropical and
subtropical regions of the Old World.

Uses: Medicinal.

P28. 齒蕨科 Acrostichaceae

齒蕨屬 *Acrostichum* L.

齒蕨 Leather Fern, Mangrove Fern

Acrostichum aureum L.

Hab.: Herb.

Loc.: Mai Po, Tai Shui Hang, Wong Nai Chung
Gap, Sha Tin, Sai Kung. On littoral mud
flat.

Dist.: Guangdong, Hainan, Yunnan. Tropics.

P29. 光葉藤蕨科

Stenochlaenaceae

光葉藤蕨屬 *Stenochlaena* J. Sm.

光葉藤蕨 Marshy Vine Fern, Palustral Vine Fern

†**Stenochlaena palustris* (Burm.) Bedd.

Hab.: Climbing fern.

Loc.: Mai Po, Sha Chau. On wet places in thin
forest.

Dist.: Hainan, Yunnan. Subtropical regions.

P30. 中國蕨科 Sinopteridaceae

粉背蕨屬 *Aleuritopteris* Fée

粉背蕨

†*Aleuritopteris farinosa* (Forsk.) Fée

Hab.: Herb.

Loc.: Tai Mo Shan. On walls.

Dist.: Fujian, Guangdong, Guangxi, Jiangxi,
Hunan, Sichuan, Yunnan. Laos,
Myanmar, India, Iran, Afghanistan,
Africa, South America.

碎米蕨屬 *Cheilosoria* Trev.

毛軸碎米蕨 Chusan Lip-fern

†*Cheilosoria chusana* (Hook.) Ching & K. H. Shing

Hab.: Herb.

Loc.: Sai Kung, Mt. Davis. In forests.

Dist.: Areas S of Qinling Range. Japan,
Philippines, Vietnam.

碎米蕨 Lace-fern, Mysure Lip-fern

Cheilosoria mysurensis (Wall. ex Hook.) Ching & K. H. Shing (*Cheilanthes* *mysurensis* auct. non Wall : Check List HK Pl. 1993.)

Hab.: Herb.

Loc.: Tai Mei Tuk, Tai Tan (Sai Kung). In
thickets and on streamsides.

Dist.: Taiwan, Fujian, Guangdong, Hainan.
Tropical Asia.

薄葉碎米蕨 Narrow-leaved Lip-fern, Thin-leaved Lip-fern

Cheilosoria tenuifolia (Burm.) Trev. (*Cheilanthes tenuifolia* Sw.)

Hab.: Herb.

Loc.: Common in Hong Kong. On hillslopes
near villages.

Dist.: Taiwan, Fujian, Guangdong, Hainan,
Guangxi, Yunnan. Tropical Asia.

烏軸蕨屬 *Dorypteryx* J. Sm.

烏軸蕨

Dorypteryx concolor (Langsd. & Fisch.) Kuhn

Hab.: Herb.

Loc.: Mirs Bay (only one specimen collected)

<p>in 1906). In forests.</p> <p>Dist.: Taiwan, Hainan, Guangdong, Guangxi. Tropics.</p> <p>隱囊蕨屬 <i>Notholaena</i> R. Br.</p> <p>隱囊蕨 <i>Hirsute Notholaena</i>, Lip-fern</p> <p>Notholaena hirsuta (Poir.) Desv. (<i>Cheilanthes hirsuta</i> Desv.)</p> <p>Hab.: Herb.</p> <p>Loc.: Shek O, Ho Chung, Lung Kwu Tan, Po Toi Islands. Along streams.</p> <p>Dist.: Taiwan, Fujian, Guangdong, Guangxi. Tropical Asia, Polynesia.</p> <p>金粉蕨屬 <i>Onychium</i> Kaulf.</p> <p>日本金粉蕨 Japanese Claw Fern</p> <p>Onychium japonicum (Thunb.) Kunze</p> <p>Hab.: Herb.</p> <p>Loc.: Aberdeen, Stanley, Lantau Island. In forests.</p> <p>Dist.: Areas S of Qinling Range. Japan, Philippines, Indonesia, Polynesia.</p> <p>Uses: Medicinal.</p>	<p>Dist.: Taiwan, Fujian, Guangdong, Hainan, Guangxi, Guizhou, Yunnan. Tropical and subtropical Asia.</p> <p>扇葉鐵線蕨 Fan-leaved Maidenhair</p> <p>Adiantum flabellulatum L.</p> <p>Hab.: Herb.</p> <p>Loc.: Common in Hong Kong. On sunny hillslopes.</p> <p>Dist.: Areas S of Changjiang River. Japan, Malaysia, Vietnam, India, Sri Lanka.</p> <p>Uses: Medicinal. Indicator of acid soil.</p> <p>南洋鐵線蕨 Walking Maidenhair Fern</p> <p>†Adiantum malesianum Ghatak</p> <p>Hab.: Herb.</p> <p>Loc.: Common in Hong Kong. In thickets.</p> <p>Dist.: S & SW China. Malaysia, Vietnam, Thailand, Myanmar, India, Sri Lanka.</p> <p>半月鐵線蕨 (菲島鐵線蕨) Philippine Maidenhair, Crescent Maidenhair</p> <p>Adiantum philippense L. (<i>Adiantum lunulatum</i> Burm.)</p> <p>Hab.: Herb.</p> <p>Loc.: Aberdeen, Shing Mun, Sha Tin. In forests.</p> <p>Dist.: Taiwan, Guangdong, Hainan, Guangxi, Guizhou, Yunnan, Sichuan. Tropical and subtropical Asia.</p>
<p>P31. 鐵線蕨科 Adiantaceae</p> <p>鐵線蕨屬 <i>Adiantum</i> L.</p> <p>鐵線蕨 Maidenhair</p> <p>Adiantum capillus-veneris L.</p> <p>Hab.: Herb.</p> <p>Loc.: Common in Hong Kong. On calcareous soil.</p> <p>Dist.: Areas S of Qinling Range. Temperate area in the world.</p> <p>Uses: Medicinal, ornamental.</p> <p>鞭葉鐵線蕨 Walking Maidenhair, Walking Fern</p> <p>Adiantum caudatum L.</p> <p>Hab.: Herb.</p> <p>Loc.: Kowloon City, Wu Shek Kok, Lantau Island. In forests.</p>	<p>P32. 水蕨科 Parkeriaceae</p> <p>水蕨屬 <i>Ceratopteris</i> Brongn.</p> <p>水蕨 Floating Fern, Water Fern</p> <p>Ceratopteris thalictroides (L.) Brongn. [<i>Ceratopteris siliquosa</i> Cop.]</p> <p>Hab.: Herb.</p> <p>Loc.: Lai Chi Chong, Sham Chung, Po Toi Islands, Ping Chau (Mirs Bay), Kam Tin, Plover Cove. In paddy field or wet</p>

places.

Dist.: Areas S of Changjiang River. Tropical and subtropical regions

Uses: Medicinal, edible.

P33. 裸子蕨科 Hemionitidaceae

粉葉蕨屬 *Pityrogramma* Link

粉葉蕨 Silver Fern

†**Pityrogramma calomelanos* (L.) Link
(*Pityrogramma tartarea* auct. non Maxon; Check List HK Pl. 1993.)

Hab.: Herb.

Loc.: North Point, Yau Ma Tei, Fanling. In valleys.

Dist.: Taiwan, Hainan, Yunnan. Tropical region.

Uses: Ornamental.

P35. 書帶蕨科 Vittariaceae

書帶蕨屬 *Vittaria* Sm.

書帶蕨 Shoe-string Fern

Vittaria flexuosa Fée (*Vittaia lineata* auct. non Sm.; Check List HK Pl. 1993.)

Hab.: Herb.

Loc.: Tai Mo Shan, Lantau Peak. On trunks or rocks.

Dist.: Areas S of Changjiang River. Japan, Vietnam, Nepal, India.

P36. 蹄蓋蕨科 Athyriaceae

短腸蕨屬 *Allantodia* R. Br.

邊生短腸蕨 Conterminal Twin-sorus Fern

†*Allantodia contermina* (Christ) Ching

Hab.: Herb.

Loc.: Hong Kong. In forests.

Dist.: S & SW China. Vietnam, Japan.

膨大短腸蕨 Widened Twin-sorus Fern

†**Allantodia dilatata* (Blume) Ching
(*Asplenium latifolium* Don; *Diplazium*

crinipes Ching)

Hab.: Herb.

Loc.: Tai Mo Shan. In forests.

Dist.: Taiwan, Fujian, S & SW China. SE Asia to Polynesia.

闊葉短腸蕨

Allantodia matthewii (Cop.) Ching
(*Asplenium matthewii* Cop.)

Hab.: Herb.

Loc.: Tai Mo Shan, Sunset Peak. In forests.

Dist.: Fujian, Guangdong, Guangxi, Yunnan. Vietnam.

江南短腸蕨

Allantodia metteniana (Miq.) Ching
[*Athyrium japonicum* auct. non (Thbg.)
Cop.: Check List HK Pl. 1993,
Diplazium mettenianum (Miq.) C. Chr.]

Hab.: Herb.

Loc.: Tai Mo Shan, Ma On Shan, Sunset Peak, Lantau Peak. In forests.

Dist.: Taiwan, Fujian, Jiangxi, Hunan, Guangdong, Hainan, Guangxi, Guizhou, Sichuan. Japan, Vietnam.

淡綠短腸蕨 Greenish Twin-sorus Fern

Allantodia virescens (Kunze) Ching

Hab.: Herb.

Loc.: Mount Parker, Tai Mo Shan. In forests.

Dist.: Areas S of Changjiang River. Vietnam, Japan.

假蹄蓋蕨屬 *Athyriopsis* Ching

假蹄蓋蕨

†*Athyriopsis japonica* (Thunb.) Ching
[*Athyrium japonicum* (Thunb.) Cop.;
Lunathyrium japonicum (Thunb.) Ching;
Lunathyrium conilii Kurata]

Hab.: Herb.

Loc.: Common in Hong Kong. In forests.

Dist.: Areas S of Changjiang River. Korea, Japan.

毛軸假蹄蓋蕨

***Athyriopsis petersenii** (Kunze) Ching
[*Athyriopsis japonica* auct. non (Thunb.) Ching : Check List HK Pl. 1993.]

Hab.: Herb.

Loc.: Victoria Peak, Wong Nai Chung, Tai Mo Shan. In forests.

Dist.: Areas S of Changjiang River. Japan, SE Asia.

菜蕨屬 **Callipteris** Bory

菜蕨 Fleshy Lady-fern

Callipteris esculenta (Retz.) J. Sm. (*Athyrium esculentum* Cop.)

Hab.: Herb.

Loc.: Aberdeen. In wet places.

Dist.: Taiwan, Fujian, Guangdong, Hainan, Guangxi, Yunnan. Asia, tropical Australia.

Uses: Edible.

雙蓋蕨屬 **Diplazium** Sw.

雙蓋蕨 Twin-sorus Fern

Diplazium donianum (Mett.) Tard.-Blot
(*Athyrium bantamense* auct. non Milde: Check List HK Pl. 1993.)

Hab.: Herb.

Loc.: Wong Nai Chung, Tung Chung. In forests.

Dist.: Taiwan, Fujian, Guangdong, Guangxi, Yunnan. India, Vietnam, Malaysia, Japan.

馬鞍山雙蓋蕨 Feathery-leaved Lady-fern

Diplazium maonense Ching [*Athyrium pinnatum* auct. non (Blanco) Cop.: Check List HK Pl. 1993.]

Hab.: Herb.

Loc.: Mount Nicholson, Lantau Peak, Ma On Shan. In forests.

Dist.: Fujian.

毛子蕨屬 **Monomelangium** Hayata

毛子蕨 Pullinger's Lady-fern

Monomelangium pullingeri (Baker) Tagawa
[*Athyrium pullingeri* (Baker) Cop.]

Hab.: Herb.

Loc.: Tai Mo Shan, Ma On Shan. In forests.

Dist.: Guangdong, Hainan, Fujian, Taiwan. SE Asia.

單葉雙蓋蕨屬 **Triblemma** (J. Sm.) Ching

單葉雙蓋蕨 Spear-leaved Lady-fern

Triblemma lancea (Thunb.) Ching [*Athyrium lanceum* (Thunb.) Milde]

Hab.: Herb.

Loc.: Common in Hong Kong. In forests.

Dist.: S, SE & SW China. SE & S Asia

裂葉雙蓋蕨 Ceylon Spleenwort

Triblemma zeylanica (Hook.) Ching
(*Athyrium zeylanica* Hook.)

Hab.: Herb.

Loc.: Caine Rd. (only one specimen collected in 1907). On rocks along streams.

Dist.: Taiwan, Fujian, Guangdong, Guangxi, Hainan, Sichuan, Hunan. Japan, SE Asia, Sri Lanka.

P38. 金星蕨科 **Thelypteridaceae**

星毛蕨屬 **Ampelopteris** Kunze

星毛蕨

†**Ampelopteris prolifera** (Retz.) Cop.

Hab.: Herb.

Loc.: Chau Tau. In thin forests, along roadside.

Dist.: S & SW China. Tropical regions of Asia, Africa.

毛蕨屬 *Cyclosorus* Link

漸尖毛蕨 Hairy Wood-fern, Acuminate Tri-vein Fern
Cyclosorus acuminatus (Houtt.) Nakai
Hab.: Herb.
Loc.: Common in Hong Kong. In forests, along roadside.
Dist.: Areas S of Qinling Range. Korea, Japan, Vietnam.

乾旱毛蕨 Dry Wood-fern, Arid Tri-vein Fern
Cyclosorus aridus (Don) Tagawa
Hab.: Herb.
Loc.: Common in Hong Kong. In forests, along roadside.
Dist.: E, S & SW China. SE Asia, India, Nepal.

齒牙毛蕨 Tapering Tri-vein Fern, Downy Woody-fern
Cyclosorus dentatus (Forsk.) Ching
Hab.: Herb.
Loc.: Tai Mo Shan. On wet places of valleys.
Dist.: Taiwan, Fujian, Jiangxi, Guangdong, Guangxi, Yunnan, Guizhou. Vietnam, Thailand, Myanmar, India, Tropical America.

異子毛蕨 Heterocarpous Tri-vein Fern
Cyclosorus heterocarpus (Blume) Ching
Hab.: Herb.
Loc.: Common in Hong Kong. In valleys.
Dist.: Guangdong, Hainan. SE Asia.

間斷毛蕨 Interrupted Tri-vein Fern
Cyclosorus interruptus (Willd.) H. Ito (*Cyclosorus gongyloides* Link)
Hab.: Herb.
Loc.: Common in Hong Kong. On wet places
Dist.: Taiwan, Fujian, Jiangxi, Guangdong, Hainan, Guangxi. S & SE Asia to

Polynesia.

寬羽毛蕨 Broad-pinna Tri-vein Fern
Cyclosorus latipinnus (Benth.) Tard.-Blot
Hab.: Herb.
Loc.: Common in Hong Kong. In valleys or along streams.
Dist.: Fujian, Guangdong, Hainan, Guangxi. SE Asia.

華南毛蕨 Wood-fern, Parasitic Tri-vein Fern
Cyclosorus parasiticus (L.) Farw. [*Christella parasitica* (L.) Leveille]
Hab.: Herb.
Loc.: Common in Hong Kong. On margins of forests and roadside.
Dist.: Taiwan, Fujian, Hunan, Guangdong, Hainan, Guangxi, Guizhou, Sichuan. Tropics.

聖蕨屬 *Dictyocline* Moore
聖蕨 Wilford's Dictyocline
Dictyocline wilfordii (Hook.) J. Sm. (*Dictyocline griffithii* auct. non Moore: Check List HK Pl. 1993.)
Hab.: Herb.
Loc.: Tai Mo Shan, Sunset Peak, Lantau Peak. In forests.
Dist.: Zhejiang, Taiwan, Fujian, Guangdong, Guangxi, Guizhou, Yunnan, Sichuan. Vietnam, Japan.

針毛蕨屬 *Macrothelypteris* (H. Ito) Ching

普通針毛蕨 Mariana Maiden Fern, Wood-fern
Macrothelypteris torresiana (Gaudich.) Ching (*Dryopteris setigera* C. Chr.)
Hab.: Herb.
Loc.: Common in Hong Kong. On wet places of valleys.

Dist.: Areas S of Changjiang River. Tropical Asia.

金星蕨屬 Parathelypteris (H. Ito) Ching

鈍頭金星蕨

†**Parathelypteris angulariloba** (Ching) Ching

Hab.: Herb.

Loc.: Victoria Peak. In forests.

Dist.: Fujian, Guangdong, Guangxi. Japan.

金星蕨

Parathelypteris glanduligera (Kunze) Ching

[*Dryopteris gracilescens* (Blume)
Kuntze; *Lastrea thelypteris* auct. non (L.)
Bory : Check List HK Pl.1993.]

Hab.: Herb.

Loc.: Victoria Peak, Mount Gough, Mount Parker, Tai Po Kau. In forests.

Dist.: Areas S of Changjiang River. Korea, Japan to India.

新月蕨屬 Pronephrium Presl

新月蕨 Rugged Pronephrium

†**Pronephrium aspera** (Presl) Shieh & Tsai
[*Pronephrium gymnopteridifrons*
(Hayata) Holttum; *Cyclosorus urophylloides* auct. non Cop.: Check List HK Pl. 1993.)

Hab.: Herb.

Loc.: Mount Parker, Tai Mo Shan, Castle Peak. In forests.

Dist.: Taiwan, Guangdong, Guangxi, Yunnan. Tropical Asia.

紅色新月蕨 Red Pronephrium

†**Pronephrium lakhimpurens** (Ros.) Holtt.

Hab.: Herb.

Loc.: Tai Mo Shan.

Dist.: Fujian, Jiangxi, Guangdong, Guangxi, Sichuan, Yunnan. India, Thailand,

Vietnam.

微紅新月蕨

Pronephrium sampsoni (Baker) Ching

[*Cyclosorus sampsoni* (Baker) Donk]

Hab.: Herb.

Loc.: Tai Mo Shan, Pat Sin Leng. In valleys.

Dist.: Fujian, Guangdong, Guangxi, Yunnan. Japan, Vietnam, Thailand.

單葉新月蕨 Simple Pronephrium

Pronephrium simplex (Hance)

Holtt.[*Cyclosorus simplex* (Hook.) Cop.]

Hab.: Herb.

Loc.: Common in Hong Kong. In valleys.

Dist.: Taiwan, Fujian, Guangdong, Hainan, Guangxi, Yunnan. Vietnam.

三羽新月蕨 Three-leaved Pronephrium

Pronephrium triphyllum (Sw.) Holtt.

[*Cyclosorus triphyllus* (Sw.) Cop.]

Hab.: Herb.

Loc.: Pak Tai To Yan. In forests.

Dist.: Taiwan, Fujian, Guangdong, Guangxi, Yunnan.

假毛蕨屬 Pseudocyclosorus Ching

溪邊假毛蕨

Pseudocyclosorus ciliatus (Wall.) Ching

(*Lastrea calcarata* Moore)

Hab.: Herb.

Loc.: Common in Hong Kong. Along streams.

Dist.: Guangdong, Hainan, Guangxi, Guizhou, Yunnan. S & SE Asia.

鱗片假毛蕨 Falcate-lobed Pseudocyclosorus

Pseudocyclosorus falciloba (Hook.) Ching

Hab.: Herb.

Loc.: Tai Mo Shan. Along streams in valleys.

Dist.: E & S China. Japan, Indochina, India.

普通假毛蕨 Common Pseudocyclosorus
Pseudocyclosorus subochthodes (Ching)
 Ching [*Pseudocyclosorus esquirolii* auct.
 non (Christ) Ching]
 Hab.: Herb.
 Loc.: Tai Mo Shan, Fo Tan. In forests.
 Dist.: Areas S of Changjiang River. Korea,
 Japan to India.

假毛蕨 False Pseudocyclosorus
 †**Pseudocyclosorus tylodes** (Kunze) Ching
 Hab.: Herb.
 Loc.: Tai Mo Shan.
 Dist.: S & SW China. S Asia to Indochina.

P39. 鐵角蕨科 Aspleniaceae

鐵角蕨屬 Asplenium L.

合生鐵角蕨 Spleenwort
Asplenium adnatum Cop.
 Hab.: Herb.
 Loc.: Tai Mo Shan (only one specimen
 collected in 1907). On streamsides.
 Dist.: Guangdong.

華南鐵角蕨 South China Spleenwort
Asplenium austrochinense Ching (*Asplenium*
wilfordii auct. non Mett. ex Kuhn.:
 Check List HK Pl. 1993.)
 Hab.: Herb.
 Loc.: Tai Mo Shan, Sunset Peak. In forests.
 Dist.: Areas S of Changjiang River. Vietnam.

齒果鐵角蕨 Lip-sorus Spleenwort
Asplenium cheilosorum Kunze ex Mett.
 Hab.: Herb.
 Loc.: Mount Parker, Tai Mo Shan. In forests.
 Dist.: Taiwan, Fujian, Hainan, Guangxi, SW
 China. Japan, SE & S Asia.

毛軸鐵角蕨 Hairy-stemmed Spleenwort

Asplenium crinicaule Hance
 Hab.: Herb.
 Loc.: Hong Kong Island, Tai Mo Shan. In
 forests.
 Dist.: Jiangxi, Guangdong, Hainan, Guangxi,
 Yunnan, Guizhou, Sichuan. S & SE Asia
 to Australia.

劍葉鐵角蕨 Sword-leaved Spleenwort
Asplenium ensiforme Wall. ex Hook. & Grev.
 (*Asplenium holosorum* Christ)
 Hab.: Herb.
 Loc.: Tai Mo Shan. In forests.
 Dist.: Taiwan, Guangdong, Guangxi, Hunan,
 Sichuan, Guizhou, Yunnan, Xizang.
 India, Nepal, Sri Lanka to Japan.

镰葉鐵角蕨 Falcate-leaved Spleenwort
Asplenium falcatum Lam. (*Asplenium*
macrophyllum auct. non Sw: Check List
 HK Pl. 1993.)
 Hab.: Herb.
 Loc.: Hong Kong Island, Lantau Island, Castle
 Peak, Chek Chau. In forests.
 Dist.: Yunnan, Guizhou, Guangdong, Hainan,
 Taiwan. S & SE Asia to Polynesia, South
 Africa.

厚葉鐵角蕨 Griffith's Spleenwort
 †**Asplenium griffithianum** Hook.
 Hab.: Herb.
 Loc.: Ng Tung Chai. In forests, on rocks along
 streams.
 Dist.: Taiwan, Fujian, Hunan, S & SW China.
 India, Myanmar, Vietnam to Japan.

大羽鐵角蕨 Wedge-shaped Spleenwort
Asplenium neolaserpitifolium Tard.-Blot &
 Ching (*Asplenium cuneatum* Lam., p.p.)
 Hab.: Herb.

Loc.: Hong Kong. In forests, on rocks along streams.	Hab.: Herb.
Dist.: Taiwan, Hainan, Yunnan. S Japan to Vietnam, India, Malaysia.	Loc.: Tai Mo Shan, Ma On Shan. Along streams.
倒掛鐵角蕨 Normal Spleenwort Asplenium normale Don	Dist.: Areas S of Changjiang River. Vietnam, Japan.
Hab.: Herb.	巢蕨屬 Neottopteris J. Sm.
Loc.: Hong Kong Island, Ma On Shan. In forests.	巢蕨 (雀巢芒) Bird-nest Fern
Dist.: Areas S of Changjiang River. S & SE Asia, Japan, Hawaii, Madagascar.	Neottopteris nidus (L.) J. Sm. (<i>Asplenium nidus</i> L.)
長葉鐵角蕨 Rue-leaved Spleenwort Asplenium prolongatum Hook.	Hab.: Herb.
Hab.: Herb.	Loc.: Mount Parker, Cape D'Aguilar, Tai Tam, Tai Pang Wan. On trees or rocks.
Loc.: Common in Hong Kong. On trees and rocks.	Dist.: Taiwan, Guangdong, Guangxi, Yunnan. Philippines, Vietnam, Malaysia.
Dist.: Areas S of Changjiang River. Japan, Korea, India, Sri Lanka, Myanmar, Vietnam, Fiji Islands.	P42. 烏毛蕨科 Blechnaceae
Uses: Ornamental.	烏毛蕨屬 Blechnum L.
假大羽鐵角蕨 Large-pinna Spleenwort † Asplenium pseudolaserpitifolium Ching	烏毛蕨 Oriental Blechnum
Hab.: Herb.	Blechnum orientale L.
Loc.: Hong Kong Island, Ng Tung Chai. In forests.	Hab.: Herb.
Dist.: Taiwan, Guangdong, Hainan, Guangxi, Yunnan. India to SE Asia.	Loc.: Common in Hong Kong. In thin forests.
半邊鐵角蕨 Unilateral Spleenwort Asplenium unilaterale Lam.	Dist.: Areas S of Changjiang River. S & SE Asia.
Hab.: Herb.	蘇鐵蕨屬 Brainea J. Sm.
Loc.: Tai Mo Shan. In forests.	蘇鐵蕨 Cycad-fern
Dist.: Areas S of Changjiang River. Tropical Asia, East Africa.	Brainea insignis (Hook.) J. Sm.
翹柄鐵角蕨 Wright's Spleenwort Asplenium wrightii Eaton ex Hook.	Hab.: Herb.
	Loc.: Common in Hong Kong. On grassy slopes or in thickets.
	Dist.: Taiwan, Fujian, Guangdong, Hainan, Guangxi, Yunnan. India, SE Asia.
	Uses: Ornamental.
	崇澍蕨屬 Chienopteris Ching
	崇澍蕨 Harland's Hain Fern
	Chienopteris harlandii (Hook.) Ching (<i>Woodwardia harlandii</i> Hook.)
	Hab.: Herb.
	Loc.: Mount Parker, Tai Mo Shan. In valleys.

Dist.: Taiwan, Fujian, Guangdong, Hainan, Guangxi, Hunan, Vietnam, Japan.

裂羽崇澍蕨 Dissected Hain Fern, Kemp's Hain Fern

†**Chieniopteris kempii** (Cop.) Ching (*Woodwardia harlandii* auct. non Hook: Check List HK Pl. 1993, p. p.)

Hab.: Herb.

Loc.: Tai Mo Shan (only one specimen collected in 1899). In forests.

Dist.: Taiwan, Fujian, Guangdong, Guangxi, Japan.

狗脊屬 **Woodwardia** Sm.

狗脊 Japanese Chain Fern

Woodwardia japonica (L. f.) J. Sm.

Hab.: Herb.

Loc.: Hong Kong Island, Tai Mo Shan. In thin forests.

Dist.: Areas S of Changjiang River. Korea, Japan.

Uses: Medicinal.

東方狗脊 Oriental Chain Fern

Woodwardia orientalis Sw.

Hab.: Herb.

Loc.: Victoria Peak, Sha Tau Kau, Tai Mo Shan, Ma On Shan. On hillslopes.

Dist.: Taiwan, Fujian, Jiangxi, Guangdong, Japan, Philippines.

芽珠狗脊 Prolific Chain Fern

†**Woodwardia prolifora** Hook. & Arn.

Hab.: Herb.

Loc.: Ho Pui Reservoir.

Dist.: E & S China. Japan.

頂芽狗脊 Unigemmate Chain Fern

†**Woodwardia unigemimata** (Makino) Nakai

Hab.: Herb.

Loc.: Hong Kong.

Dist.: Areas S of Qinling Range. Japan, S & SE Asia.

P45. 鱗毛蕨科 **Dryopteridaceae**

芒蕨屬 **Arachniodes** Blume

中華芒蕨

†**Arachniodes chinensis** (Ros.) Ching

Hab.: Herb.

Loc.: Victoria Peak, Ng Tung Chai. In forests.

Dist.: Areas S of Changjiang River. Vietnam, Japan.

芒蕨

Arachniodes exilis (Hance) Ching [*Rumohra aristata* auct. non (Forst) Ching: Check List HK Pl. 1993.]

Hab.: Herb.

Loc.: Ng Tung Chai. In forests.

Dist.: Areas S of Changjiang River.

黑鱗芒蕨

Arachniodes nigrospinosa (Ching) Ching

Hab.: Herb.

Loc.: Tai Mo Shan. In forests.

Dist.: S China.

斜方芒蕨

***Arachniodes rhomboidea** (Wall. ex Mett.) Ching [*Rumohra amabilis* auct. non (Blume) Ching: Check List HK Pl. 1993.]

Hab.: Herb.

Loc.: Hong Kong. In forests.

Dist.: Areas S of Changjiang River. Japan, Myanmar, India, Nepal.

鞭葉蕨屬 **Cyrtomidictyum** Ching

單葉鞭葉蕨

Cyrtomidictyum basipinnatum (Baker) Ching (*Aspidium basipinnatum* Baker)

Hab.: Herb.

Loc.: Mount Nicholson (only one specimen collected in 1906). In forests.

Dist.: S China.

貫衆屬 **Cyrtomium** Presl

鑊羽貫衆 Falcate Holly Fern

Cyrtomium balansae (Christ) C. Chr.
[*Phanerophlebia vittata* Cop.]

Hab.: Herb.

Loc.: Ma On Shan, Lantau Island. In forests.

Dist.: Areas S of Changjiang River. Vietnam, Japan.

全緣貫衆 Asian Holly Fern

Cyrtomium falcatum (L. f.) Presl
[*Phanerophlebia falcata* (L. f.) Cop.]

Hab.: Herb.

Loc.: Shek O, Tap Mun Chau, Ninepin Group. On rocks.

Dist.: Shandong, Jiangsu, Zhejiang, Taiwan, Fujian, Guangdong. Japan.

Uses: Medicinal.

鱗毛蕨屬 **Dryopteris** Adans.

闊鱗鱗毛蕨 Male-fern

Dryopteris championii (Benth.) C. Chr. ex Ching [*Dryopteris filix-mas* auct. non (L.) Schott: Check List HK Plants. 1993.]

Hab.: Herb.

Loc.: Victoria Peak, Mount Parker. In forests.

Dist.: Areas S of Changjiang River. Japan.

異蓋鱗毛蕨

Dryopteris decipiens (Hook.) Kuntze

Hab.: Herb.

Loc.: Tai Mo Shan. In forests.

Dist.: Fujian, Jiangxi, Guangdong, Guangxi, Guizhou. Japan.

黑足鱗毛蕨 Autumn Fern

Dryopteris fuscipes C. Chr. [*Dryopteris erythrosora* auct. non (Eaton) Kuntze: Check List HK Pl. 1993.]

Hab.: Herb.

Loc.: Tai Mo Shan.

Dist.: Areas S of Changjiang River. Japan, Indochina.

柄葉鱗毛蕨

Dryopteris podophylla (Hook.) Kuntze

Hab.: Herb.

Loc.: Common in Hong Kong. In forests.

Dist.: Fujian, Guangdong.

稀疏鱗毛蕨

Dryopteris sparsa (Don) Kuntze

Hab.: Herb.

Loc.: Tai Mo Shan, Ma On Shan. In forests.

Dist.: Areas S of Changjiang River. India, Vietnam, Japan.

嶺南鱗毛蕨

†**Dryopteris tenuicula** Matthew & Christ

Hab.: Herb.

Loc.: No precise locality.

Dist.: Guangdong.

變異鱗毛蕨

Dryopteris varia (L.) Kuntze [*Polystichum varium* (L.) Presl]

Hab.: Herb.

Loc.: Common in Hong Kong. In forests.

Dist.: Areas S of Changjiang River. Japan.

耳蕨屬 **Polystichum** Roth

華南耳蕨 Green-grey Shield Fern

†**Polystichum eximium** (Mett. ex Kuhn) C. Chr.

Hab.: Herb.

Loc.: Tai Mo Shan. On hillslopes.

Dist.: S China, Taiwan, Vietnam, Japan.

P46. 叉蕨科 Asplidiaceae

沙皮蕨屬 *Hemigramma* Christ

沙皮蕨

Hemigramma decurrens (Hook.) Cop.

Hab.: Herb.

Loc.: Hong Kong Island, Nam Chung, Sha Tau Kau. In forests.

Dist.: Taiwan, Fujian, Guangdong, Hainan, Yunnan, Vietnam, Japan.

黃腺蕨屬 *Pleocnemia* Presl

有蓋黃腺蕨

Pleocnemia conjugata (Blume) Presl [*Tectaria leuzeana* auct. non (Gaudich.)

Cop.: Check List HK Pl. 1993, p. p.]

Hab.: Herb.

Loc.: Mount Parker, Ho Chung. In forests.

Dist.: SE Asia, Myanmar, India.

黃腺蕨

Pleocnemia winitii Holtt. [*Tectaria leuzeana* auct. non (Gaudich.) Cop.: Check List HK Pl. 1993, p. p.]

Hab.: Herb.

Loc.: Mount Gough, Pat Sin Leng. In forests.

Dist.: Taiwan, Fujian, Guangdong, Hainan, Guangxi, Yunnan, Vietnam, Thailand, India.

地耳蕨屬 *Quercifilix* Cop.

地耳蕨 *Ceylon Quercifilix*

Quercifilix zeylanica (Houtt.) Cop.

Hab.: Herb.

Loc.: Common in Hong Kong. In forests.

Dist.: Taiwan, Fujian, Guangdong, Guangxi, Yunnan, Guizhou, Vietnam, Malaysia, India, Sri Lanka.

叉蕨屬 *Tectaria* Cav.

下延叉蕨 Decurrent Halberd Fern

†*Tectaria decurrens* (Presl) Cop.

Hab.: Herb.

Loc.: Mount Parker. In forests.

Dist.: Taiwan, S China, Yunnan, India, Sikkim, Myanmar, Vietnam, Philippines, Japan.

條裂叉蕨

Tectaria phaeocaulis (Ros.) C. Chr. (*Tectaria laciniata* Ching)

Hab.: Herb.

Loc.: New Territories, Mount Parker, Tai Mo Shan, Tate's Cairn, Ma On Shan. In forests.

Dist.: Taiwan, Fujian, Guangdong, Guangxi, Vietnam, Japan.

叉蕨

Tectaria subtriphylla (Hook. & Arn.) Cop.

Hab.: Herb.

Loc.: Common in Hong Kong. In forests.

Dist.: Taiwan, S & SW China, India, Sri Lanka, Myanmar, Vietnam, Indonesia.

P47. 實蕨科 Bolbitidaceae

實蕨屬 *Bolbitis* Schott

華南實蕨

Bolbitis subcordata (Cop.) Ching [*Bolbitis quoyana* auct. non (Gaut.) Ching: Check List HK Pl. 1993.]

Hab.: Herb.

Loc.: Tai Mo Shan, Tai Po Kau. Along streams in forests.

Dist.: E & S China, Vietnam, Japan.

刺蕨屬 *Egenolfia* Schott

刺蕨

Egenolfia appendiculata (Willd.) J. Sm.

Hab.: Herb.

Loc.: Tai Mo Shan, Mount Parker. In forests.

Dist.: Taiwan, Guangdong, Guangxi, Yunnan, SE Asia, Japan, India.

中華刺蕨

Egenolfia sinensis (Baker) Maxon

Hab.: Herb.

Loc.: Sunset Peak, Tai Po Kau. In forests.

Dist.: Yunnan, Guizhou. S & SE Asia.

P48. 藤蕨科 Lomariopsidaceae

藤蕨屬 Lomagramma J. Sm.

網藤蕨

Lomagramma matthewii (Ching) Houtt.

Hab.: Climbing herb.

Loc.: Ng Tung Chai. On cliffs near waterfall.

Dist.: Fujian, Guangdong, Yunnan.

P49. 舌蕨科 Elaphoglossaceae

舌蕨屬 Elaphoglossum Schott

華南舌蕨

Elaphoglossum yoshinagae (Yatabe) Makino

(*Elaphoglossum austrosinicum* Metthew
& Christ)

Hab.: Herb.

Loc.: Tai Mo Shan, Sunset Peak, Lantau Peak.
On rocks in forest.

Dist.: E & S China, Hunan, Guizhou. Japan.

F50. 腎蕨科 Nephrolepidaceae

腎蕨屬 Nephrolepis Schott

腎蕨 Tuberous Sword Fern

Nephrolepis auriculata (L.) Trimem

(*Nephrolepis cordifolia* auct. non C.
Presl: Check List HK Pl. 1993.)

Hab.: Herb.

Loc.: Common in Hong Kong. In forests.

Dist.: E, S & SE China. Tropical and
subtropical regions

Uses: Medicinal, ornamental.

長葉腎蕨 Broad Sword-fern

Nephrolepis biserrata (Sw.) Schott

Hab.: Herb.

Loc.: Hong Kong Island. In forests.

Dist.: Taiwan, Guangdong, Hainan, Yunnan.

Japan, Vietnam, India to Malaysia.

毛葉腎蕨 Rough Sword Fern

Nephrolepis hirsutula (Forst.) Presl

[*Nephrolepis exaltata* auct. non (L.)
Schott: Check List HK Pl. 1993.]

Hab.: Herb.

Loc.: Hong Kong Island. In forests.

Dist.: Taiwan, S China, Yunnan. India, Vietnam,
Malaysia.

P51. 條蕨科 Oleandraceae

條蕨屬 Oleandra Cav.

圓基條蕨

Oleandra intermedia Ching (*Oleandra*

cumingii auct. non J. Sm.: Check List
HK Pl. 1993.)

Hab.: Herb.

Loc.: Mount Parker (only two specimens
collected in 1906 to 1907). On cliffs.

Dist.: Guangdong, Guizhou, Yunnan.

P52. 骨碎補科 Davalliaceae

骨碎補屬 Davallia Sm.

華南骨碎補 South China Hare's-foot Fern

†**Davallia austrosinica** Ching

Hab.: Herb.

Loc.: Hong Kong.

Dist.: Guangdong.

大葉骨碎補 Large Hare's-foot Fern

Davallia formosana Hayata

Hab.: Herb.

Loc.: Victoria Peak, Tai Mo Shan. In valleys.

Dist.: Taiwan, Fujian, Guangdong, Hainan,
Guangxi, Yunnan. Vietnam, Cambodia.

陰石蕨屬 **Humata** Cav.

陰石蕨 Repent Humate

Humata repens (L. f.) Diels

Hab.: Herb.

Loc.: Victoria Peak, Aberdeen, Lantau Island.
On trees or rocks.

Dist.: E, S & SW China.

Uses: Medicinal, ornamental.

圓蓋陰石蕨 Bear's-foot Fern

Humata tyermannii Moore

Hab.: Herb.

Loc.: Wong Nai Chung, Tai Tam, Shek Kong.
On trees or rocks.

Dist.: S, E & SW China. Vietnam, Laos.

Uses: Medicinal.

P56. 水龍骨科 **Polypodiaceae**

線蕨屬 **Colysis** Presl

線蕨(蛇眼草) Snake's-eye Fern

Colysis elliptica (Thunb.) Ching

Hab.: Herb.

Loc.: Common in Hong Kong. In forests.

Dist.: Areas S of Changjiang River. Vietnam,
Japan.

斷線蕨 Interrupted Colysis

Colysis hemionitidea (Wall. ex Mett.) Presl

Hab.: Herb.

Loc.: Mount Parker, Tai Mo Shan. In forests.

Dist.: Taiwan, Fujian, Guangdong, Guangxi,
Guizhou, Yunnan. Japan, Philippines,
Vietnam, Myanmar, India.

寬羽線蕨 Broad Colysis

Colysis pothifolia (D. Don) C. Presl [*Colysis elliptica* var. *pothifolia* (D. Don) Ching]

Loc.: Sai Kung, Tai Mo Shan, Tai To Yan. In
forests.

Dist.: Areas S of Changjiang River. S & SE

Asia.

褐葉線蕨 Brown Colysis

Colysis wrightii (Hook.) Ching

Hab.: Herb.

Loc.: Pak Tai To Yan, Tai Om, Sheung Wo
Hang, Bride's Pool. On streamsides.

Dist.: Taiwan, Fujian, Jiangxi, Guangdong,
Guangxi. Vietnam, Japan.

伏石蕨屬 **Lemmaphyllum** Presl

伏石蕨

Lemmaphyllum microphyllum Presl

Hab.: Herb.

Loc.: Common in Hong Kong. On trees or
rocks.

Dist.: Areas S of Changjiang River. Korea,
Japan, Vietnam.

骨牌蕨屬 **Lepidogrammitis** Ching

披針骨牌蕨

†**Lepidogrammitis diversa** (Ros.) Ching

Hab.: Herb.

Loc.: Sunset Peak. On rocks in forest.

Dist.: Anhui, Zhejiang, Jiangxi, Fujian, Hunan,
Guangdong, Guangxi, Guizhou.

Uses: Medicinal.

骨牌蕨

***Lepidogrammitis rostrata** (Bedd.) Ching

[*Lemmaphyllum subrostratum* auct. non
(C. Chr.) Ching: Check List HK Pl.
1993.]

Hab.: Herb.

Loc.: Ma On Shan. On rocks in forest.

Dist.: Taiwan, Guangdong, Guangxi, Yunnan,
Guizhou. Vietnam, Myanmar, India.

瓦韋屬 *Lepisorus* (J. Sm.) Ching

粵瓦韋

†*Lepisorus obscure-venulosus* (Hayata) Ching

Hab.: Herb.

Loc.: Tai Mo Shan, Lantau Peak. On trunks.

Dist.: S & E China. Japan.

瓦韋 Thunberg's *Lepisorus*

Lepisorus thunbergianus (Kaulf.) Ching

Hab.: Herb.

Loc.: Mount Parker, New Territories. On trees or rocks.

Dist.: Areas S of Changjiang River. Japan, Korea.

Uses: Medicinal.

星蕨屬 *Microsorium* Link

攀援星蕨

Microsorium buergerianum (Miq.) Ching

Hab.: Herb.

Loc.: Common in Hong Kong. On trees or rocks.

Dist.: Areas S of Changjiang River. Vietnam, Japan.

羽裂星蕨

Microsorium insigne (Blume) Cop.

[*Microsorium dilatatum* (Bedd.) Sledge;
Microsorium hancockii Ching]

Hab.: Herb.

Loc.: Aberdeen, Mount Gough, Bride's Pool, Tai Mo Shan, Mui Tsz Lam. On rocks in forest.

Dist.: Taiwan, Fujian, Guangdong, Guangxi, Yunnan, Guizhou, Sichuan. India, Malaysia, Vietnam, Japan.

江南星蕨

Microsorium fortunei (Moore) Ching

Hab.: Herb.

Loc.: Castle Peak. On trunks or rocks.

Dist.: Areas S of Changjiang River. Vietnam, Japan.

Uses: Medicinal, ornamental.

有翅星蕨

Microsorium pteropus (Blume) Cop.

Hab.: Herb.

Loc.: Aberdeen, Shing Mun, Fanling. On streamsides.

Dist.: Taiwan, Fujian, Guangdong, Guangxi, Guizhou, Yunnan. India, Vietnam.

星蕨

Microsorium punctatum (L.) Cop.

Hab.: Herb.

Loc.: Tai Mo Shan, Castle Peak, Tan Chuk Hang, Chek Chau. On trunks.

Dist.: Taiwan, Guangdong, Hainan, Yunnan, Guizhou. Tropical Asia, Africa.

顯脈星蕨

Microsorium zippelii (Blume) Ching

Hab.: Herb.

Loc.: Tai Mo Shan. On cliffs along stream.

Dist.: S China, Yunnan, Guizhou. Philippines, Sumatra, Malaysia, N India.

瘤蕨屬 *Phymatodes* Presl

多羽瘤蕨

Phymatodes longissima (Blume) J. Sm.

Hab.: Herb.

Loc.: Tai Mo Shan, Sai Kung, Wu Kau Teng, Lamma Island, Hei Ling Chau. On streamsides and in thin forests.

Dist.: Taiwan, Hainan. Tropical Asia.

瘤蕨

Phymatodes scolopendria (Burm. f.) Ching

Hab.: Herb.

Loc.: North Point, Tai Hang, Tap Mun Chau, Centre Island, Chek Chau, Round Island, Sha Chau. On rocks in forest.

Dist.: Taiwan, Hainan, Guangdong, Yunnan. Tropical Asia, Africa.

石韋屬 **Pyrrosia** Mirbel

貼生石韋 Tongue-fern, Close-growing Pyrrosia

Pyrrosia adnascens (Sw.) Ching [*Pyrrosia lanceolata* (L.) Farw.]

Hab.: Herb.

Loc.: Common in Hong Kong. On rocks or trees.

Dist.: Taiwan, Fujian, Guangdong, Guangxi, Yunnan. Vietnam, India, Malaysia.

Uses: Medicinal.

石韋 Japanese Felt Fern

Pyrrosia lingua (Thunb.) Farw.

Hab.: Herb.

Loc.: Common in Hong Kong. On trees or rocks.

Dist.: Areas S of Changjiang River. Vietnam, Japan.

Uses: Medicinal, ornamental.

F57. 檵蕨科 **Drynariaceae**

崖薑屬 **Pseudodrynaria** C. Chr.

崖薑 Rock-ginger Fern

Pseudodrynaria coronans (Wall. ex Mett.) Ching

Hab.: Large herb.

Loc.: Common in Hong Kong. In forests.

Dist.: Taiwan, Fujian, Guangdong, Guangxi, Guizhou, Yunnan. India, Vietnam, Malaysia.

Uses: Medicinal, ornamental.

P59. 禾葉蕨科 **Grammitidaceae**

禾葉蕨屬 **Grammitis** Sw.

紅毛禾葉蕨

Grammitis hirtella (Blume) Tuyame

Hab.: Herb.

Loc.: Tai Mo Shan. On rocks along streams.

Dist.: Guangdong. Tropical Asia.

兩廣禾葉蕨

†***Grammitis lasiosora** (Blume) Ching

Hab.: Herb.

Loc.: Tai Mo Shan, Sunset Peak. In forests.

Dist.: Fujian, Guangdong, Guangxi, Jiangxi, Hunan. India, Malaysia.

P60. 劍蕨科 **Loxogrammaceae**

劍蕨屬 **Loxogramme** (Blume) Presl

柳葉劍蕨 Narrow-leaved Sword Fern

Loxogramme salicifolia (Makino) Makino
(*Loxogramme lanceolata* C. Presl)

Hab.: Herb.

Loc.: Tai Mo Shan, Lantau Peak. On trees or rocks.

Dist.: Areas S of Changjiang River. Korea, Japan, Vietnam.

P61. 蘋科 **Marsileaceae**

蘋屬 **Marsilea** L.

蘋 Water Shamrock

Marsilea quadrifolia L.

Hab.: Herb.

Loc.: Sha Tin, Mai Po, Pui O. In paddy fields, ponds and ditches.

Dist.: Areas S of Changjiang River, N China. Temperate and tropical regions.

Uses: Medicinal.

P62. 槐葉蘋科 **Salviniaceae**

槐葉蘋屬 **Salvinia** Adans.

槐葉蘋 Water Spangles

Salvinia natans (L.) All.

Hab.: Herb.

Loc.: Mau Ping, Hang Ha Po . In paddy fields,
ponds and ditches.

Dist.: Widely distributed in China. (except
NW areas). India, Vietnam, Japan,
Europe.

Uses: Medicinal.

P63. 滿江紅科 **Azollaceae**

滿江紅屬 **Azolla** Lam.

滿江紅 Mosquito Fern

Azolla imbricata (Roxb.) Nakai (*Azolla pinnata* R. Br. var. *africana* auct. non Baker: Check List HK Pl. 1993.)

Hab.: Herb.

Loc.: Tai Po, Sha Tau Kau, Tan Chuk Hang. In
paddy fields, ponds and ditches.

Dist.: Areas S of Changjiang River. Korea,
Japan.

Uses: Medicinal, forage, green manure.